

**SECOND SEMESTER B.C.A. DEGREE EXAMINATION, MAY 2014**

(U.G.-CCSS)

Complementary Course

CA 2C 04—NUMERICAL METHODS IN C

Time : Three Hours

Maximum : 30 Weightage

**Part A (Objective Type Questions)***Answer all twelve questions.*

1. The numbers in the computer word can be stored in two forms. Which are they ?
2. Define the inherent error.
3. When we can say that is a root of the equation  $f(x) = 0$ .
4. Define the central difference operator  $\delta$ .
5. Write Newton's forward difference approximation of  $O(h^2)$ .
6. What is the formula to find  $\int_a^b f(x) dx$  using Simpson's rule ?

Fill in the blanks :

7. To avoid the difficulty of keeping every number less than 1 in magnitude during computation, most computers use \_\_\_\_\_ representation for a real number.
8. Bisection method is based on the repeated application of the \_\_\_\_\_ theorem.
9. In Gauss-Jordan elimination method the coefficient matrix is reduced to a \_\_\_\_\_ matrix.
10. If there are  $n + 1$  distinct points  $x_0 < x_1 < x_2 < \dots < x_n$ , then the problem of Lagrange and Newton interpolation for the continuous function  $f(x)$  on  $[a, b]$  is to obtain  $p(x)$  satisfying the conditions \_\_\_\_\_
11. The **Hermite** interpolating polynomial interpolates not only the function  $f(x)$  but also its \_\_\_\_\_ at a given set of tabular points.
12. The general problem of numerical integration is to find an approximate value of the integral  $I = \int_a^b w(x) f(x) dx$  where  $w(x) > 0$  in  $[a, b]$ .

(12 x  $\frac{1}{4}$  = 3 weightage)**Turn over**

**Part B (Short Answer Questions)**

*Answer all nine questions.*

13. Find the decimal number corresponding to the binary number  $(111 \cdot 011)_2$ .
14. Construct the difference table for the sequence of values  $f(x) = (0, 0, 0, \epsilon, 0, 0, 0)$ .
15. Solve the equations  $x + y = 2$  and  $2x + 3y = 5$  by Gauss-Jordan method.
16. State intermediate value theorem.
17. Evaluate  $\int_0^4 e^x dx$  by Simpson's '1/3' rule using the data  $e = 2.72, e^2 = 7.39, e^3 = 20.09$  and  $e^4 = 54.60$ .
18. Perform 2 iterations of the bisection method to obtain a real root of the equation  $x^3 - x - 11 = 0$ .
19. Solve  $\frac{dy}{dx} = 1 - y, y(0) = 0$  using Euler's method. Find  $y$  at  $x = 0.1$ .
20. Find the  $n^{\text{th}}$  difference of  $e^x$ .
21. Show that  $\mu = [1 + 8^2 / 4]^{1/2}$ .

(9 x 1 = 9 weightage)

**Part C (Short Essay Questions)**

*Answer any five questions.*

22. Apply Cramer's rule to solve the equations,  $3x + y + 2z = 3, 2x - 3y - z = 3$  and  $x + 2y + z = 4$ .
23. Solve the following system of equations using Gaussian elimination method  $x + y + z = 9, 2x - 3y + 4z = 13$  and  $3x + 4y + 5z = 40$ .
24. Construct Newton's forward interpolation polynomial for the following data :
 

$x :$	4	6	8	10
$y :$	1	3	8	16
25. Evaluate  $\int_0^{10} \frac{dx}{1 + x^2}$  by using Trapezoidal rule.

26. Using Taylor's method, find  $y(0.1)$  from  $\frac{dy}{dx} + 2xy = 1, y_0 = 0$ .
27. Evaluate  $\sqrt{12}$  to four places of decimals by ~~Newton-Raphson~~ method.
28. The equation  $8x^3 - 12x^2 - 2x + 3 = 0$  has 3 real roots in the interval  $[-2, 3]$ . Find the intervals each of unit length containing each one of these roots.

(5 x 2 = 10 weightage)

**Part D (Essay Questions)***Answer any two questions.*

29. (a) Write the Lagrange's interpolation formula.  
 (b) Use Lagrange's formula to find the value of  $y$  at  $x = 6$  from the following data :

$x :$	3	7	9	10
	168	120	72	63

30. (a) Find  $y'(x)$  given :

$x$	:	0	1	2	3	4
$y(x)$	:	1	1	15	40	85

- (b) The population of a certain town is shown in the following table :

Year $x$	:	1931	1941	1951	1961	1971
Population in 1961 $y$	:	40.62	60.80	79.95	103.56	132.65

31. (a) What is the relation between ~~Runge-Kutta~~ method and modified ~~Euler's~~ method.  
 (b) Use ~~Runge-Kutta~~ method of the fourth order to find  $y(0.1)$  given that :

$$\frac{dy}{dx} = \frac{1}{x+y}, y(0) = 1.$$

(2 x 4 = 8 weightage)