

FOURTH SEMESTER M.Sc. DEGREE EXAMINATION, JUNE 2016

(CUCSS)

Mathematics

MT 4C 15—FUNCTIONAL ANALYSIS – II

Time : Three Hours

Maximum : 36 Weightage

Part A

*Answer all questions.**Each question carries 1 weightage.*

1. Show that if F is a bijective closed map on a normed space then F^{-1} is also a closed map.
2. Let X be a Banach space and $A \in \mathcal{BL}(X)$. Show that if $\|A^n\| \leq \frac{1}{n}$, then $\|A\| \leq 1$.
3. Let X be a normed space and $A \in \mathcal{BL}(X)$ be invertible. Show that $\liminf_{n \rightarrow \infty} \|A^n\|^{1/n} > 0$.
4. Let Y be a subspace of a normed space X . For $x \in X$, let $F(x) = \frac{x}{\|x\|}$. Show that F is a surjective linear map from X to \bar{Y} such that $\|F(x)\| = 1$ for all $x \in X$.
5. Let X be a Banach space. Show that if X is reflexive then it remains reflexive in any equivalent norm.
6. Let $M = \text{diag}(k_1, k_2, \dots)$ and X be the sequence space. Show that if $k_n \rightarrow 0$ as $n \rightarrow \infty$, then M defines a map in $\mathcal{CL}(X)$.
7. State Riesz representation theorem.
8. Let H be a Hilbert space and (x_n) be a sequence in H . Show that $x_n \rightharpoonup x$ in H if $\langle x_n, y \rangle \rightarrow \langle x, y \rangle$ uniformly for $y \in H$ with $\|y\| \leq 1$.
9. Let X be a non-zero Banach space and $P \in \mathcal{BL}(X)$ be a projection. Show that if $0 \neq P$ then $\sigma(P) = \{0, 1\}$.

Turn c

10. Let H be the Hilbert space \mathbb{C}^2 and $A : H \rightarrow H$ be defined by :

$$A(x(1), x(2)) = (2x(1), x(2)) \text{ for } (x(1), x(2)) \in H. \text{ Determine } A^*.$$

11. Let H be a Hilbert space. Show that if (A_n) is a sequence of unitary operators on H , and $A \in \mathcal{BL}(H)$ be such that $\|A_n - A\| \rightarrow 0$ as $n \rightarrow \infty$, then A is unitary.

12. Let H be a Hilbert space and $A \in \mathcal{BL}(H)$ be self-adjoint. Show that $A^2 \geq 0$ and $\|A\| \leq \|A\|$.

13. Show that if x_1 and x_2 are eigen vectors of a normal operator on a Hilbert space H corresponding to distinct eigen values, then $x_1 \perp x_2$.

14. Let H be a non-zero Hilbert space and $A \in \mathcal{BL}(H)$. Show that :

$$\|A\| = \sup \{ \|Ax\| : \|x\| = 1 \}$$

(14 x 1 = 14 weightage)

Part B

Answer any seven questions.
Each question carries 2 weightage.

15. Let X and Y be Banach spaces and $F \in \mathcal{BL}(X, Y)$. Show that $R(F)$ is linearly homeomorphic to $X/Z(F)$ iff $R(F)$ is closed in Y .

16. Let X be a Banach space and $A \in \mathcal{BL}(X)$. Show that A is invertible iff A is bounded below and the range of A is dense in X .

17. Let X be the sequence space l^2 and $A : X \rightarrow X$ be defined by :

$$A(x) = (0, x(1), \frac{x(2)}{2}, \frac{x(3)}{3}, \dots) \text{ for } x \in X. \text{ Show that } \sigma_p(A) = \emptyset \text{ and } \sigma_w(A) = \{0\} = \sigma(A).$$

18. Let X and Y be normed spaces and $F \in \mathcal{BL}(X, Y)$. Show that $\|F\| = \|J_X F\|$ and $F'' J_X = J_Y F$, where J_X and J_Y are the Canonical embedding of X and Y into X'' and Y'' respectively.

19. Let X be a reflexive normed space show that X is separable iff X' is separable.
20. Let X and Y be Banach spaces and $F : X \rightarrow Y$ be linear. Show that F is continuous and of finite rank iff F is a compact map and $R(F)$ is closed in Y .
21. Let H be a Hilbert space and $f \in H'$. Show that if $g \in H'$ and $g(x) = f(x)$ for some non-zero $x \in Z(f)$, then $g = f$.
22. Let H be a Hilbert space and $A \in \mathcal{BL}(H)$ be self-adjoint. Show that
- $$\|A\| = \sup\{|\langle Ax, x \rangle| : x \in H, \|x\| \leq 1\}.$$
23. Let H be a Hilbert space and $A \in \mathcal{BL}(H)$. Show that $\sigma_n(A) \subset \overline{w(A)}$ and $\sigma(A)$ is contained in the closure of $w(A)$.
24. Let H be a Hilbert space and $A \in \mathcal{BL}(H)$. Show that A is compact iff A^*A is compact.

(7 x 2 = 14 weightage)

Part C*Answer any two questions.**Each question carries 4 weightage.*

25. State and prove closed graph theorem.
26. Let F be a finite dimensional subspace of a Hilbert space H . Show that $H = F + F^\perp$ and $F^{\perp\perp} = F$.
27. Let H be a Hilbert space and $A \in \mathcal{BL}(H)$. Show that $R(A) = H$ iff A^* is bounded below, and $R(A^*) = H$ iff A is bounded below.
28. Let A be a non-zero compact self adjoint operator on a Hilbert space H over K . Show that there exist a finite or infinite sequence (s_n) of non-zero real numbers with $|s_1| \geq |s_2| \geq |s_3| \geq \dots$ and an orthonormal set $\{u_1, u_2, \dots\}$ in H . Such that $A(x) = \sum s_n \langle x, u_n \rangle u_n, x \in H$.

(2 x 4 = 8 weightage)